

Possible Participation of Lithium in Fleischmann-Pons Reaction is Testable

P. Frodl¹, O. E. Rössler², M. Hoffmann², and F. Wahl³

¹ Institute for Physics, University of Mainz, D-6500 Mainz.

² Institute for Physical and Theoretical Chemistry, University of Tübingen, D-7400 Tübingen, FRG.


³ Institute for Theoretical Physics, University of Tübingen, D-7400 Tübingen, FRG.

Z. Naturforsch. **45a**, 757–758 (1990); received May 2, 1989

If the amount of heat reported by Fleischmann and Pons to occur in their electrochemical “fusion” experiment is assumed to be produced entirely by the “clean” nuclear reaction ${}^6\text{Li} + {}^2\text{D} \rightarrow 2\text{}^4\text{He}$ (22.4 MeV), then the concentration of ${}^6\text{Li}$ in the reaction fluid should during a period of 120 hours go down by a measurable amount, namely between 3.4 and 0.25 percent dependent on whether the natural isotopic mixture of (less likely) pure ${}^6\text{Li}$ was used in the experiment.

Fleischmann and Pons [1] recently reported on an electrochemical experiment in which “too much” heat was produced to be accounted for by chemical reaction. Nevertheless, only a very small amount of gamma radiation, neutron flux production and tritium production was observed – by a factor of 10^9 too small, in fact, if one assumes ${}^2\text{D} + {}^2\text{D}$ reactions to account for the heat. The authors therefore hypothesized that an “unknown” nuclear reaction may be participating.

While the details of the experiments have yet to be confirmed by other groups – so far only inconclusive and contradictory results are available – the question is interesting enough to be taken seriously in its own right. Recently, Jones [2] in a different context discussed as an especially clean nuclear reaction the example of



Among the various Li + D-reactions we concentrate on (1); for this reaction is “appealing because there are no neutrons in the final state” [2]. As it happens, also the radiation production is exceedingly low. The reaction (1) is of potential interest because lithium has been used by Fleischmann and Pons as a 0.1 molar solution within the heavy water electrolyzed, in contrast to the second recently claimed electrochemically induced “cold fusion” by Jones et al. [4], who had not

Reprint requests to Dr. P. Frodl, Institut für Physik, D-6500 Mainz, FRG.

reported an excessively high production of caloric energy. Assuming the reaction (1) to be responsible for the major part of the apparent heat, the measured neutrons and gammas might stem from subordinate ${}^2\text{D} + {}^2\text{D}$ reactions.

Of course, a mechanism for the deuterons either colliding with each other or colliding with other light nuclei inside the palladium is still pending. The hypothesis that the positive deuterons might be “delocalized” within the palladium crystal among the positive palladium ions, with the consequence that a Bose condensation (with the fermion subparticles forming a degenerate Fermi gas) might possibly form, was recently proposed [5]. This condensation might potentially be responsible for a “surreptitious” mutual approach up to the minimum tunneling distance required for the nuclear reaction in question to occur. This, of course, is only a speculation at the time being.

At any rate, there is evidence that lithium can be taken up into the palladium crystal much like hydrogen and the noble gases [6]. Therefore there is the possibility that, if for some reason the deuterons could come close to the lithium, the above reaction might take place. Although the question would have to be answered how competing ${}^2\text{D} + {}^6\text{Li}$ reactions leading to byproducts like neutrons, protons or gammas could be suppressed, it is perhaps of interest to see whether the reaction named (1) can *in principle* account for the observed energy production as well as the cleanness reported.

Fleischmann and Pons [1] reported that $14.6 \cdot 10^{23}$ deuterium atoms were present in their experiment using the rod-shaped electrode, which lasted for 120 hours. This yields a volume employed of 21.6 ml. A 0.1 molar solution of LiOD in heavy water then contains $1.3 \cdot 10^{21}$ lithium ions. To obtain an energy output of 26.8 watts, $7.5 \cdot 10^{12}$ reactions (1) would be needed per second. In 120 hours, about $3.2 \cdot 10^{18}$ fusion events would occur.

This means that, if all lithium ions present originally were ${}^6\text{Li}$, about 0.25 percent would have had to be used up at the end of the experiment. If, on the other hand, natural lithium (7.3 percent ${}^6\text{Li}$, the rest ${}^7\text{Li}$) has been used as likely, 3.4 percent would have had to be used up. *Both* rates of decrease are readily measurable with current methods.

To conclude, we propose to monitor the ${}^6\text{Li}$ concentration in future runs of the Fleischmann-Pons exper-

0932-0784 / 90 / 0500-0757 \$ 01.30/0. – Please order a reprint rather than making your own copy.



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

iment. In addition the amount of the ^4He produced could be measured too. Because of probable energy transfers to the Pd-lattice, a measurement of the α -radiation has to be complemented by an analysis of the Pd electrode with respect to embedded α 's.

Acknowledgements

P. Frodl would like to thank C. Tappe for stimulating conversations. Discussions with A. Faessler are also gratefully acknowledged.

Note added in proof: After our paper was submitted, two analogous ideas have been reported [7, 8]. That lithium might play a crucial role is indicated by some results of J. Appleby [9]. However, our proposal is not contingent on the ultimate outcome of the Fleischmann-Pons debate, but provides another possibility to falsify (or – perhaps – corroborate) their conjecture of nuclear reactions.

- [1] M. Fleischmann and S. Pons, *J. Electroanal. Chem.* **261**, 301 (1989).
- [2] S. E. Jones, *Nature London* **321**, 127 (1986).
- [3] F. Ajzenberg-Selove, *Nucl. Phys. A* **413**, 1 (1984).
- [4] S. E. Jones et al., *Nature London* **338**, 737 (1989).
- [5] O. E. Rössler, J. Becker, M. Hoffmann, and W. Nadler, *Z. Naturforsch.* **44a**, 329 (1989).
- [6] C. Tappe, personal communication.
- [7] J. S. Cohen and J. D. Davies, *Nature London* **338**, 705 (1989).
- [8] B. U. R. Sundqvist et al., *Physica Scripta* **40**, 303 (1989).
- [9] Cited in: R. Pool, *Science* **244**, 1039 (1989).