

The Space Group of $(\text{TMM-TTF})^{2+}(\text{AuCl}_4^-)_2$ [TMM-TTF = Tetra(methylthio)tetra- thiofulvalene]

Peter G. Jones*

Institut für Anorganische Chemie der Universität,
Tammannstraße 4, 3400 Göttingen

Z. Naturforsch. **44b**, 243–244 (1989);
received October 10, 1988

Tetra(methylthio)tetrathiofulvalene

The space group of the title compound is probably $C2/m$ rather than the previously reported $C2$.

During the preparation of a review article on the crystal structures of gold compounds, it was noticed that the structure of $(\text{TMM-TTF})^{2+}(\text{AuCl}_4^-)_2$, presented in space group $C2$ with cell constants $a = 14.799$, $b = 11.513$, $c = 8.034$ Å, $\beta = 97.05^\circ$ [1], displayed several unusual features. The atoms Au, Cl(2), Cl(4) and C(1) lay in or near the planes $y = 0$ or $1/2$; other atoms occurred in pairs with similar x and z coordinates, and y coordinates summing to 0 or 1; and some chemically equivalent bond lengths differed appreciably (Au–Cl 2.247–2.309 Å, S–C(1) 1.573, 1.787 Å). All these features could be ex-

plained if the true space group were $C2/m$, *i.e.* if a crystallographic centre of symmetry had been overlooked [2].

The deposited structure factors were obtained from the Fachinformationszentrum Energie, Physik, Mathematik, 7514 Eggenstein-Leopoldshafen 2 (deposition number CSD 52689). Anisotropic refinement in $C2/m$ proceeded smoothly to an R value of 0.032 for 1511 reflections, 70 parameters (*cf.* 0.028 for 1581 reflections, 140 parameters in $C2$; only 1512 reflections were present in the deposited material and one was clearly in error). Atoms Au, Cl(2) and Cl(3) (in the new numbering scheme, see Tables I and II) were fixed in the mirror planes. The weighting scheme was $w = \sigma^{-2}(F)$, with σ values calculated by the method of Cruickshank [3] (no σ values had been deposited). H atoms were not included. The program system was SHELX-76, locally modified by its author Prof. G. M. Sheldrick.

The new refinement leads to more regular geometry, with Au–Cl 2.255–2.275 Å, C(1)–S(1) 1.684 Å, and to much lower e.s.d.'s [2]. The descrip-

Table I. Atomic coordinates ($\times 10^4$) and equivalent isotropic displacement parameters ($\text{Å}^2 \times 10^3$).

	x	y	z	$U(\text{eq})^*$
Au	1448.9(2)	0	1147.5(4)	44(1)
Cl(1)	1498(2)	1974(2)	1252(3)	72(1)
Cl(2)	2495(2)	0	– 689(4)	67(1)
Cl(3)	350(3)	0	2860(5)	83(1)
S(1)	470(1)	3756(2)	8307(2)	50(1)
S(2)	1376(1)	3584(2)	5204(2)	60(1)
C(1)	211(6)	5000	9244(9)	40(2)
C(2)	959(4)	4390(6)	6722(7)	42(2)
C(3)	1360(6)	2137(9)	6067(10)	72(3)

* Equivalent isotropic U defined as one third of the trace of the orthogonalized U_{ij} tensor.

Table II. Bond lengths (Å) and angles ($^\circ$).

Au–Cl(1)	2.275(2)	Au–Cl(2)	2.266(3)
Au–Cl(3)	2.255(4)	S(1)–C(1)	1.684(5)
S(1)–C(2)	1.704(6)	S(2)–C(2)	1.707(7)
S(2)–C(3)	1.806(10)	C(1)–C(1a)	1.433(16)
Cl(1)–Au–Cl(2)	90.2(1)	Cl(1)–Au–Cl(3)	90.0(1)
Cl(2)–Au–Cl(3)	177.0(1)	Cl(1)–Au–Cl(1b)	174.8(1)
C(1)–S(1)–C(2)	96.4(3)	C(2)–S(2)–C(3)	101.8(4)
S(1)–C(1)–S(1c)	116.5(5)	S(1)–C(1)–C(1a)	121.7(2)
S(1)–C(2)–S(2)	121.7(4)	S(1)–C(2)–C(2c)	115.4(2)
S(2)–C(2)–C(2c)	122.9(2)		

Symmetry operators: (a) $-x, y, 2-z$; (b) $x, -y, z$; (c) $x, 1-y, z$




Fig. 1. Thermal ellipsoid plot (50% level) of the title compound, showing the numbering of the asymmetric unit in $C2/m$.

* Current address: Institut für Anorganische und Analytische Chemie der Technischen Universität, Hagenring 30, D-3300 Braunschweig

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen
0932–0776/89/0200–0243/\$ 01.00/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition “no derivative works”). This is to allow reuse in the area of future scientific usage.

tion in $C2/m$ thus seems preferable. The anion possesses exact m symmetry and the cation exact $2/m$ symmetry.

Observed and calculated structure factors in $C2/m$ and anisotropic thermal parameters have been deposited as above. Any request for this material

should quote the new reference number CSD 53456 and a full literature citation.

I thank the Fonds der Chemischen Industrie for financial support and the authors of the original report [1] for helpful correspondence.

-
- [1] K. Brunn, H. Endres, and J. Weiss, *Z. Naturforsch.* **43b**, 224 (1988).
[2] D. W. G. Cruickshank and W. S. McDonald, *Acta Crystallogr.* **23**, 9 (1967).

- [3] D. W. G. Cruickshank, D. F. Pilling, A. Bujosa, F. M. Lovell, and M. R. Truter, "Computing Methods and the Phase Problem in X-Ray Crystal Analysis", Oxford, Pergamon (1961).