

# Absolute Configuration of Chiral 1,3,4,5-Tetrahydro-2*H*-1,5-benzodiazepin-2-ones

Mashooda Hasan<sup>a</sup>, Tanveer Ahmad Nizami<sup>b</sup>, Abdul Malik<sup>c</sup>, Claus Ohlendorf<sup>d</sup>,  
Wolfgang Hiller<sup>d</sup>, Wolfgang Voelter<sup>b,\*</sup>

<sup>a</sup> Department of Chemistry, Quaid-i-Azam University, Islamabad, Pakistan

<sup>b</sup> Abteilung für Physikalische Biochemie, Physiologisch-chemisches Institut der  
Universität Tübingen, Hoppe-Seyler-Straße 4, D-72076 Tübingen, Germany

<sup>c</sup> On leave from the H. E. J. Research Institute of Chemistry University of Karachi,  
Karachi-75270, Pakistan

<sup>d</sup> Anorganisch-chemisches Institut der Technischen Universität München,  
Lichtenbergstraße-4, D-85747 Garching, Germany

Z. Naturforsch. **53b**, 485–487 (1998); received January 19, 1998

Chiral 1,5-Benzodiazepin-2-ones, Crystal Structure, Absolute Configuration

X-ray studies are carried out to correlate the Cotton effects in CD spectra with the absolute configuration of a series of previously reported chiral 1,3,4,5-tetrahydro-2*H*-1,5-benzodiazepin-2-ones.

\* Reprint requests to Prof. Dr. W. Voelter.