

X-Ray Structure Analysis of *iso*-Sildenafil (*iso*-Viagra) [1]

Mustafa M. El-Abadelah^{a,*}, Salim S. Sabri^a, Monther A. Khanfar^a,
Wolfgang Voelter^{b,*}, and Cäcilia Maichle-Mössmer^c

^a Chemistry Department, Faculty of Science, University of Jordan, Amman, Jordan

^b Abteilung für Physikalische Biochemie, Physiologisch-chemisches Institut der
Universität Tübingen, Hoppe-Seyler-Straße 4, D-72076 Tübingen, Germany

^c Institut für Anorganische Chemie auf der Morgenstelle 18, Universität Tübingen,
D-72076 Tübingen, Germany. Fax: (+49) 7071-293348.
E-mail: wolfgang.voelter@uni-tuebingen.de

* Reprint requests to Prof. M. M. El-Abadelah or to Prof. W. Voelter

Z. Naturforsch. **54b**, 1323–1326 (1999); received December 28, 1998

X-Ray Data, *iso*-Viagra, Cyclic Guanosine Monophosphate, Viagra, Phosphodiesterase

For structure-activity reasons, an isomeric compound of Viagra, methylated at N-2 at the pyroazolo[4,3-d]pyrimidine-7-one nucleus, was synthesized and its X-ray structure performed. The crystallographic data show that the phenyl and condensed pyrazolo-pyrimidone ring systems are not coplanar to each other.