Cyclic Oxalylation of Primary *N*-Substituted Anthranilamides: 1*H*-Benzo[*e*][1,4]diazepine-2,3,5(4*H*)-triones and 11a-Chloro-benzo[*e*]oxazolo[3,2-*a*][1,4]diazepine-2,3,5,11(10*H*,11a*H*)-tetraones

Maria Anna Köllner and Detlef Geffken

Institute of Pharmacy, University of Hamburg, Bundesstraße 45, 20146 Hamburg, Germany

Reprint requests to Prof. Dr. Detlef Geffken. Fax: +49 40 428383477.
E-mail: geffken@chemie.uni-hamburg.de

Z. Naturforsch. **2010**, *65b,* 1155 – 1160; received April 9, 2010

Dedicated to Professor Gerwalt Zinner

In dependence on the molar ratio, the reaction of primary anthranilamides **3** with oxalyl chloride produced 1*H*-benzo[*e*][1,4]diazepine-2,3,5(4*H*)-triones **4** or 10-substituted 11a-chloro-benzo[*e*]-oxazolo[3,2-*a*][1,4]diazepine-2,3,5,11(10*H*, 11a*H*)-tetraones **5**, the structures of which were unambiguously proven by X-ray diffraction analysis.

Key words: Anthranilamides, Cyclization, Fused Heterocycles, Oxalyl Chloride, Acylation