

## Enzymatic Synthesis of Sinapoyl-L-Malate from 1-Sinapoylglucose and L-Malate by a Protein Preparation from *Raphanus sativus* Cotyledons

Norbert Tkotz † and Dieter Strack

Botanisches Institut der Universität zu Köln, Gyrhofstr. 15,  
D-5000 Köln 41

Z. Naturforsch. **35 c**, 835–837 (1980);  
received June 26, 1980

*Raphanus*, Brassicaceae, Phenylpropanoid Metabolism,  
Transacylation, Sinapoyltransferase

Protein preparations from cotyledons of *Raphanus sativus* transfer the sinapoyl moiety of 1-sinapoylglucose to L-malate to form sinapoyl-L-malate. To our knowledge this is the first time that this type of reaction has been found to be involved in phenylpropanoid depside formation. Most acylation reactions have so far been demonstrated to depend on acyl-CoAs and only one example is known describing 1-O-acyl sugar derivatives as acyl donors.

### Introduction

Aromatic acids in higher plants are involved in a great diversity of esterification reactions and it is well established that their formation requires in most cases a CoA-activated acid [1]. Glucose esters are formed from free acids and UDP-glucose [2–5].

Glucose esters have been proposed as possibly essential intermediates in hydroxycinnamic acid metabolism [6–8]. Schlepphorst and Barz [9] have postulated from a biochemical point of view the formation of an ester from an energy-rich acyl glucose as the acyl donor in a transacylation reaction. This hypothetical reaction has recently been proven by Michalczuk and Bandurski [10], who demonstrated *in vitro* the enzymatic synthesis of IAA-myoinositol by the transfer of the IAA moiety from IAA-glucose to myo-inositol, catalyzed by an IAA-transferase. These authors demonstrated also, that the preceding reaction is the enzymatic synthesis of IAA-glucose from IAA and UDP-glucose.

The present report describes the same biochemical mechanism in the phenylpropanoid metabolism of *Raphanus sativus* cotyledons for the formation of sinapoylmalate [11]. During early stages of *Raphanus* germination, the seed constituent sinapine (sinapoylcholine) is rapidly degraded [8], catalyzed by a

specific sinapine esterase [12]. The freed sinapic acid is reesterified to 1-sinapoylglucose [13] via UDP-glucose [5]. In later stages of *Raphanus* germination, the amount of 1-sinapoylglucose in the cotyledons decreases, whereas sinapoylmalate is formed. Protein, extracted from this tissue, catalyzes the transfer of the sinapoyl moiety from 1-sinapoylglucose to L-malate. Analogous to the biosynthesis of IAA-myoinositol, the activity catalyzing the formation of sinapoylmalate might be classified as 1-sinapoylglucose:L-malate sinapoyltransferase.

### Materials and Methods

Plant material and culture conditions are described elsewhere [8].

Protein preparation was obtained from 6–7 day-old *Raphanus* cotyledons as described in ref. [5]. The enzymatic activity was prepared from 30 to 60%  $(\text{NH}_4)_2\text{SO}_4$  precipitable fraction followed by chromatography on Sephadex G-25. The incubation mixture in experiment *a* contained 20 µl protein solution, 1 µmol malate (L- or D-form), and 0.5 µmol 1-sinapoylglucose in 220 µl potassium phosphate buffer (100 mM, pH 6.5). The formation of sinapoylmalate (at room temperature) was determined by TLC and HPLC. In experiment *b* a glucosyltransferase assay [5], containing sinapic acid and UDP-glucose, was mixed with the protein preparation used in experiment *a*, to which L-malate was added at the same concentration. 1-Sinapoylglucose and authentic sinapoylmalate were isolated from *Raphanus* seedlings as described in ref. [8]. D- and L-malate were purchased from Roth, Karlsruhe.

Chromatography is described in Table I and Fig. 1.

### Results and Discussion

When protein preparations from 6–7 day-old *Raphanus* cotyledons were incubated with 1-sinapoylglucose and L-malate, a product was formed which could not be distinguished from authentic sinapoylmalate (Table I). After 18 h of incubation time, ca. 40% of 1-sinapoylglucose was converted to sinapoylmalate, whereas 60% was hydrolyzed to give free sinapic acid. The latter indicates the metabolic activity of 1-sinapoylglucose [9]. Fig. 1 shows HPLC analyses of the assay at  $t_0$  and after 18 h incubation.

Reprint requests to Dr. D. Strack.

0341-0382/80/0900-0835 \$ 01.00/0



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht:  
Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.


Table I. Spectral properties and chromatographic behaviour of sinapoylmalate [11] and a product formed by a *Raphanus* protein preparation, which was incubated with L-sinapoylglucose and L-malate.

Compound	Spectral analysis $\lambda_{\max}$ in $\text{CH}_3\text{OH}$ , nm		TLC $R_F \times 100$ <sup>b</sup>				Polyamide-CC [8] Elutable with			$t_R$ [sec]
	alone	+ NaOH <sup>a</sup>	CAW	BAW	TAW	$\text{H}_2\text{O}$	$\text{H}_2\text{O}$	$\text{CH}_3\text{OH}$	0.025% $\text{NH}_4\text{OH}$ in $\text{CH}_3\text{OH}$	
Reaction product	324	385	82	79	24	90	—	—	+	1603
Sinapoylmalate	322	384	82	79	24	90	—	—	+	1602

<sup>a</sup> 0.1 ml 0.25 N aqueous NaOH to 0.9 ml methanolic solution of the compound.

<sup>b</sup> Determined on microcrystalline cellulose (Avicel): CAW, chloroform-acetic acid-water (3:2, water saturated); BAW, n-butanol-acetic acid-water (6:1:2); TAW, toluene-acetic acid-water (2:1, water saturated).

<sup>c</sup> See Fig. 1.



Protein preparations which were not subjected to  $(\text{NH}_4)_2\text{SO}_4$  fractionation showed ca. 85% hydrolysis.

In assays without protein or containing heat denatured protein no product was formed. D-malate was unable to accept sinapic acid in the sinapoyltransferase reaction.

In an enzyme assay to determine glucosyltransferase activity [5], to which a protein preparation from 6–7 day-old cotyledons and L-malate were added (experiment b), we observed the conjugation of ca. 4% sinapic acid with L-malate, obviously via the energy-rich acyl glucoside intermediate.

We might deduce from the results described in this paper and in ref. [5] and [12] that in cotyledons of *Raphanus sativus* the following reaction sequence leads to the formation of sinapoyl-L-malate:

- (1) sinapoylcholine  
 $\rightarrow$  sinapic acid + choline
- (2) sinapic acid + UDP-glucose  
 $\rightarrow$  1-sinapoylglucose + UDP
- (3) 1-sinapoylglucose + L-malate  
 $\rightarrow$  sinapoyl-L-malate + glucose.

Fig. 1. HPLC analyses of an enzyme assay with a protein preparation from *Raphanus* cotyledons, 1-sinapoylglucose, and L-malate at  $t_0$  and after 18 h incubation. The chromatograph used was Spectra Physics (Santa Clara, Cal., USA) and is described in ref. [14]. The applied column (250×4 mm) was prepacked with LiChrosorb RP-8 (5 µm) (Merck, Darmstadt). Separation was accomplished by gradient elution: in 40 min linear from solvent A (0.5% acetic acid in water) to 50% solvent B (acetonitrile) in A+B. The flow-rate was 1 ml/min and detection was achieved at 330 nm with a Schoeffel SF 770 UV-VIS detector (Kratos Instr., Trappenkamp). 20 µl of incubation medium was directly injected without pretreatment.

The alternative biochemical mechanism for the synthesis of sinapoylmalate, proceeding *via* a sinapoyl-coenzyme-A-ester intermediate, which would be dependent on the reversibility of the *in vivo* formation of 1-sinapoylglucose by a glucosyltransferase [5], seems to be unlikely since all attempts failed so far to demonstrate any sinapoyl CoA-ligase activity.

Experiments are underway to purify the sinapoyltransferase activity for enzyme characterization, and

further studies have to show the specificity of the transacylase. It should be of interest to study other systems for their capability of an enzymatically catalyzed acylation by a 1-O-acyl sugar.

#### Acknowledgement

The support by the Deutsche Forschungsgemeinschaft is gratefully acknowledged.

- [1] M. H. Zenk, Biochemistry of Plant Phenolics, (T. Swain, J. B. Harborne, and C. F. Van Sumere, eds.), Vol. 12, 139, Plenum Press, New York 1979.
- [2] J. J. Corner and T. Swain, *Nature* **207**, 634 (1965).
- [3] J.-J. Macheix, C. R. Acad. Sc. **284**, D-33 (1977).
- [4] G. Jacobelli, M. J. Tabone, and D. Tabone, *Bull. Soc. Chim. Biol.* **40**, 955 (1958).
- [5] D. Strack, *Z. Naturforsch.* **35 c**, 204 (1980).
- [6] J. B. Harborne and J. J. Corner, *Biochem. J.* **81**, 242 (1961).
- [7] M. Kojima and I. Uritani, *Plant Cell Physiol.* **13**, 1075 (1972).
- [8] D. Strack, *Z. Pflanzenphysiol.* **84**, 139 (1977).
- [9] R. Schlepphorst and W. Barz, *Planta Med.* **36**, 333 (1979).
- [10] L. Michalczuk and R. S. Bandurski, *Biochem. Biophys. Res. Commun.* **93**, 588 (1980).
- [11] M. Linscheid, D. Wendisch, and D. Strack, *Z. Naturforsch.*, in press.
- [12] G. Nurmann and D. Strack, *Z. Naturforsch.* **34 c**, 715 (1979).
- [13] D. Strack, N. Tkotz, and M. Klug, *Z. Pflanzenphysiol.* **89**, 343 (1978).
- [14] D. Strack, G. B. Feige, and R. Kroll, *Z. Naturforsch.* **34 c**, 679 (1979).