

## Cephalic Volatiles Identified in Workers of *Mourella caerulea*, a Rare Stingless Bee Recently Rediscovered in Southern Brazil

Dieter Wittmann

Zoologisches Institut der Universität,  
Auf der Morgenstelle 28, D-7400 Tübingen,  
Bundesrepublik Deutschland

Gunther Lübke

Wittko Francke

Institut für Organische Chemie und Biochemie  
der Universität, Martin-Luther-King-Platz 6,  
D-2000 Hamburg 13, Bundesrepublik Deutschland

Z. Naturforsch. **44c**, 325–326 (1989);  
received September 1, 1988

*Mourella caerulea*, Stingless Bee, Cephalic Volatiles

From workers of *Mourella caerulea*, an extremely rare and primitive species of neotropical stingless bees, 101 cephalic volatiles were identified by GC/MS analyses.

The stingless bee species *Mourella caerulea* was first described in 1900 by Friese [1]. Since then only a few worker specimens were recorded in subtropical Brazil. The nest architecture, males and the queen were unknown until several of the subterranean nests were recently discovered during a four years survey of the bees of Rio Grande do Sul, the southernmost State of Brazil [2]. Apart from our interest in chemical communication in stingless bees [3–8], we are aware of the potential information which an analysis of cephalic volatiles, together with comparative studies on morphology and nest architecture [9] as well as on intranidal behavior [10] will contribute to a proper taxonomic collocation of *M. caerulea*, a species which has often been considered one of the most primitive living stingless bees of the neotropics.

A nest collected at Canguçu, Rio Grande do Sul, Brazil, was transferred to the laboratory. Heads of adult worker bees of different age, mostly field bees, were cut off and extracted in pentane. Extracts were analyzed using the coupled gas chromatographic/mass spectroscopic system Varian MAT 311 A [3]. The relative proportions of the identified volatile components were determined and classified as: + = < 0.05%, ++ = 0.05–0.5%, +++ = 0.5–20%, ++++ = > 20%. The following compounds were found:

Reprint requests to Dr. D. Wittmann.

Verlag der Zeitschrift für Naturforschung, D-7400 Tübingen  
0341–0382/89/0300–0325 \$ 01.30/0

Hydrocarbons	
Pentadecane	+
Heptadecane	+
7-Heptadecene	+
9-Heptadecene	+
Heptadecadiene	+
Octadecane	+
Nonadecane	+
7-Nonadecene	+
9-Nonadecene	+
Eicosane	+
Heneicosane	+++
7-Heneicosene	+
9-Heneicosene	+
Docosane	+
Tricosane	+++
7-Tricosene	+
9-Tricosene	+
2-Methyltricosane	++
Pentacosane	+++
7-Pentacosene	+
9-Pentacosene	+
Heptacosane	+++
Heptacosene*	+++
Nonacosane	+++
Nonacosene**	+++
Hentriacontane	+++
Hentriacontene***	+++
Alcohols	
2-Pentanol	+
2-Hexanol	+
2-Heptanol	++
2-Octanol	+
2-Nonanol	+
2-Pentadecanol	+
Ketones	
2-Hexanone	+
2-Heptanone	+++
2-Octanone	++
2-Nonanone	+
2-Decanone	+
2-Undecanone	+
2-Tridecanone	++
2-Pentadecanone	+
2-Heptadecanone	+
6-Methylhept-5-en-2-one	+
Aldehydes	
Hexadecanal	+
Octadecanal	++
Carboxylic acids	
Acetic acid	+
2-Methylpropionic acid	+
Butanoic acid	+
3-Methylbutanoic acid	++
Pentanoic acid	+
Hexanoic acid	+
Heptanoic acid	+
Octanoic acid	+
Nonanoic acid	+
Decanoic acid	++



Dieses Werk wurde im Jahr 2013 vom Verlag Zeitschrift für Naturforschung in Zusammenarbeit mit der Max-Planck-Gesellschaft zur Förderung der Wissenschaften e.V. digitalisiert und unter folgender Lizenz veröffentlicht: Creative Commons Namensnennung-Keine Bearbeitung 3.0 Deutschland Lizenz.

Zum 01.01.2015 ist eine Anpassung der Lizenzbedingungen (Entfall der Creative Commons Lizenzbedingung „Keine Bearbeitung“) beabsichtigt, um eine Nachnutzung auch im Rahmen zukünftiger wissenschaftlicher Nutzungsformen zu ermöglichen.

This work has been digitalized and published in 2013 by Verlag Zeitschrift für Naturforschung in cooperation with the Max Planck Society for the Advancement of Science under a Creative Commons Attribution-NoDerivs 3.0 Germany License.

On 01.01.2015 it is planned to change the License Conditions (the removal of the Creative Commons License condition "no derivative works"). This is to allow reuse in the area of future scientific usage.

Dodecanoic acid	+++
Tetradecanoic acid	+++
Hexadecanoic acid	+++
Octadecanoic acid	+++
9-Octadecenoic acid	++++
Octadecadienoic acid	++++
Octadecatrienoic acid	++++
Benzoic acid	+
Phenylacetic acid	+
Esters	
Ethyl octanoate	+
Ethyl nonanoate	+
Ethyl decanoate	+
Ethyl dodecanoate	+
Methyl tetradecanoate	+
Ethyl tetradecanoate	+
Methyl hexadecanoate	+
Ethyl hexadecanoate	++
Ethyl 7(?)-hexadecanoate	+
Methyl heptadecanoate	+
Methyl octadecanoate	+
Ethyl octadecanoate	++
Methyl 9-octadecenoate	++
Ethyl 9-octadecenoate	+++
Methyl octadecadienoate	++
Ethyl octadecadienoate	+++
Methyl octadecatrienoate	+
Ethyl octadecatrienoate	+++
2-Heptyl acetate	+
2-Heptyl octanoate	+
2-Heptyl nonanoate	+

Mixtures of positional isomers: \* = 7, 9, 10, 13; \*\* = 7, 9–14; \*\*\* = 7, 9–15.

Up to now pheromonal bouquets of only a few species of stingless bees have been studied [11]. Recently we described the occurrence of 70 volatile compounds in cephalic secretions of *Scaptotrigona postica* workers [7]. The analyzed head extracts con-

tain secretions of the mandibular glands, however, we cannot exclude that some of the compounds are extracted from the hypopharyngeal and labial glands and/or the cuticula. The list of cephalic volatiles presented here shows that *Mourella caerulea* has a complex bouquet comparable to other stingless bee species: It is made up by uneven numbered, saturated and unsaturated hydrocarbons, methylcarbinols, respective methylketones, carboxylic acids and esters. Only very few compounds show a branched carbon skeleton. The ester fraction is particularly interesting, since it contains, with three exceptions, only methyl and ethyl esters which are known as typical volatile compounds of blossoms. Only three esters of 2-heptanol were recorded. The wax ester type, represented by compounds formed from alcohols and carboxylic acids of comparable chain length, is entirely absent. Esters of this type were repeatedly recorded in many bee species and represent characteristic constituents of their cephalic secretions [12, 13].

The monospecific genus *Mourella* is probably related to *Plebeia* [14]. A comparative study of cephalic secretions in *Plebeia* species seems desirable before any chemotaxonomic conclusions can be drawn. The list of identified *Mourella* volatiles, on the other hand, can be used for bioassays in order to decode the significance of pheromonal signals in the social behavior of these bees.

#### Acknowledgements

This study was supported by a Feodor Lynen scholarship from the Alexander von Humboldt-Foundation, the DFG and the Fonds of the Chemical Industry.

- [1] H. Friese, Jur. Természetráji Füzetek **23**, 381 (1900).
- [2] D. Wittmann and M. Hoffmann, *Iheringia* (1989), in press.
- [3] W. Francke, W. Schröder, E. Engels, and W. Engels, *Z. Naturforsch.* **38c**, 1066 (1983).
- [4] W. Engels, *Adv. Invertebr. Reprod.* **4**, 285 (1986).
- [5] W. Engels, *Mem. Inst. Oswaldo Cruz* **82**, III, 35 (1987).
- [6] E. Engels, W. Engels, W. Schröder, and W. Francke, *J. Chem. Ecol.* **13**, 371 (1987).
- [7] D. Wittmann, R. Radtke, W. Francke, and W. Schröder, *Abstr. IUSI Bayreuth*, 26 (1987).
- [8] R. Radtke, D. Wittmann, W. Francke, and W. Schröder, *Abstr. IUSI Bayreuth*, 27 (1987).
- [9] J. M. F. Camargo and D. Wittmann, *Stud. neotrop. Fauna and Environ.*, in preparation.
- [10] D. Wittmann, L. Bego, and R. Zucchi, in preparation.
- [11] M. Blum, Academic Press, N.Y. 1981.
- [12] W. Francke, W. Schröder, G. Bergström, and J. Tengö, *Nova Acta Regia Societates Scientiarum Upsaliensis*, Ser. V: C, III, 127–136 (1984).
- [13] W. Francke, *Adv. Invertebr. Reprod.* **4**, 327 (1986).
- [14] H. F. Schwarz, *Bull. Am. Mus. Nat. Hist.* **90**, 500 (1948).