

**Caffeoyl Quinic and Tartaric Acids and
Flavonoids from *Lapsana communis*
L. subsp. communis (Asteraceae)**

D. Fontanel^a, C. Galtier^b, C. Viel^a and
A. Gueiffier^b

Laboratoires de Pharmacognosie^a et Chimie Thérapeu-
tique^b, Faculté des Sciences Pharmaceutiques, Université
de Tours, 31 Avenue Monge, 37 200 Tours, France

Z. Naturforsch. **53c**, 1090–1092 (1998);
received August 5/October 7, 1998

Lapsana communis *L. subsp. communis*, Lactuceae,
Asteraceae, Caffeic Acid Esters, Flavonoids

Six hydroxycinnamic acids : caffeic acid, chlorogenic acid, 3,5-O-dicaffeoylquinic acid 2-O-caffeoyltartaric acid (caftaric acid) and 2,3-O-dicaffeoyltartaric acid (chicoric acid) have been isolated from *Lapsana communis* *L. subsp. communis* aerial parts. Among flavonoids, only isoquercitrin, luteolin and luteolin-7-O- β -glucuronide were identified. Except for chlorogenic acid, these compounds represent the first report in *Lapsana communis*. Chicoric acid is the major phenylpropanoic constituent in this plant.

Reprint requests to D. Fontanei. Fax: +33/(0)2 47 36 72 39.