

Strobilurin N and Two Metabolites Related to Chorismic Acid from the Fruit Bodies of *Mycena crocata* (Agaricales)

Malcolm S. Buchanan^a, Wolfgang Steglich^{a,*} and Timm Anke^b

^a Institut für Organische Chemie der Universität, Butenandtstr. 5–13 (Haus F), D-81377 München, Germany. Fax: +49–89–2180–7756. E-mail: wos@cup.uni-muenchen.de

^b Lehrbereich Biotechnologie der Universität, Paul-Ehrlich-Straße 23, D-67663 Kaiserslautern, Germany

* Author for correspondence and reprint requests

Z. Naturforsch. **54c**, 463–468 (1999); received March 18/April 12, 1999

Mycena crocata, Toadstool, Strobilurins, Chorismic Acid Derivatives, Fungicide

Strobilurin N (**1**), a new member of the strobilurin family of antibiotics, has been isolated from the toadstool *Mycena crocata*, together with dehydrochorismic acid lactone (**5**) and 3-[1-carboxyvinyl)oxy]benzoic acid (**6**), which are closely related to chorismic acid (**10**). Their structures were determined mainly by ¹H and ¹³C NMR spectroscopy. Interestingly, strobilurin N is the first strobilurin without antifungal activity.