

Flavonoids and Terpenoids from *Luma gayana* (Barn.) Burret

Gerald A. Wächter^a, Aree Wangmaneerat^a,
Kay M. Caple^a, Gloria Montenegro^b and
Barbara N. Timmermann^{a,*}

^a Department of Pharmacology and Toxicology, College
of Pharmacy, The University of Arizona, Tucson,
AZ 85721, U. S. A. Telefax ++01-520-626-4063.
E-mail: btimmer@pharmacy.arizona.edu

^b Departamento de Ecología, Pontificia Universidad
Católica de Chile, Casilla 114-D, Santiago, Chile

* Author for correspondence and reprint requests

Z. Naturforsch. **54c**, 1140–1142 (1999);
received July 15/August 6, 1999

Luma gayana (Barn.) Burret, Myrtaceae, Flavonoids,
Terpenoids, Alkyl *p*-Coumarates

The flavonoids 5-hydroxy-7-methoxyflavanone, 6,8-dimethyl-5,7-dihydroxyflavanone and 2',4'-dihydroxy-6'-methoxy-3',5'-dimethylchalcone, a mixture of alkyl esters of *p*-coumaric acid, the triterpenoids oleanolic acid and maslinic acid, the monoterpene 1 α ,2 β ,4 β -trihydroxy-*p*-menthane, the sesquiterpene clovandiol and β -sitosterol were isolated from the aerial parts of *Luma gayana* (Barn.) Burret. This is the first report on the chemistry of this species.