

## Coumarin Glycosides from the Mosses *Dendrologotrichum dendroides* and *Dawsonia superba*\*

Juliane Raubuch, Hans Geiger and  
Hans Dietmar Zinsmeister\*\*

Fachrichtung Botanik, Universität des Saarlandes,  
D-66041 Saarbrücken

Z. Naturforsch. **54c**, 456–457 (1999);  
received February 9/March 2, 1999

Dedicated to Professor Robert Hegnauer, Leiden,  
on the occasion of his 80<sup>th</sup> birthday

Mosses, Polytrichales, *Dendrologotrichum dendroides*,  
*Dawsonia superba*, 5, 7, 8-Trihydroxy-coumarin-5-  
glucoside derivatives

*Dendrologotrichum dendroides* and *Dawsonia superba*, which belong to two exclusively Gondwanalandic genera of the Polytrichales, accumulate the same type of coumarins as species of the cosmopolitan genera of that order. The structures of the isolated compounds, one of which is a new natural product, were elucidated spectroscopically.

### Introduction

A chromatographic study by Jung (1993) suggested that species of the Polytrichales are chemically characterized by the accumulation of tri- and tetrahydroxycoumarin derivatives. In the case of representative species from the almost cosmopolitan genera *Atrichum*, *Oligotrichum*, *Pogonatum*, and *Polytrichum* this has been verified by the isolation of pure compounds and elucidation of their structures (Jung *et al.*, 1994 and Raubuch, 1998). In this communication we shall report the isolation and identification of trihydroxycoumarin derivatives from *Dendrologotrichum dendroides* and *Dawsonia superba*, representatives of two polytrichalean genera with an exclusive Gondwanalandic distribution.

Reprint requests to Prof. Zinsmeister.  
Fax: 0681/812825

\* Publication No. 134 of the Arbeitskreis Chemie und Biologie der Moose.

\*\* Present address: Fasanenweg 2, 66133 Saarbrücken, Germany.

### Results and Discussion

*D. dendroides* yielded as its main coumarin 5, 8-dihydroxy-7-methoxycoumarin-5- $\beta$ -D-glucopyranoside (**1**). This compound occurs also in all Polytrichaceae that have been studied so far (Jung *et al.*, 1994 and Raubuch, 1998). It was identified by its NMR spectra and a chromatographic comparison with authentic material. Further constituents of this moss could not be identified, because we had only a very small sample at our disposition.

For the same reason *D. superba* yielded also only its main coumarin in a quantity that was sufficient for NMR-analysis. It turned out to be a new compound – 5, 7, 8-trihydroxycoumarin-5, 7-di- $\beta$ -D-glucoside (**2**). This structure was deduced by comparison of its <sup>1</sup>H- and <sup>13</sup>C-NMR spectra with published data of related compounds and some NOE-experiments (*c.f.* Jung *et al.*, 1994 and 1995). The multiplicities and chemical shifts of its <sup>1</sup>H as well as <sup>13</sup>C resonances in the “aromatic range” indicated that **2** is a 5, 7, 8-trihydroxycoumarin derivative. The signals in the “sugar range” revealed that **2** contains two  $\beta$ -glucopyranoses, which are both linked directly to the aglycone, NOE interactions between H-6 and the anomeric protons of both glucoses proved that these sugars are linked to O-5 and O-7. Thus **2** is 3, 7, 8-trihydroxycoumarin-5, 7-di- $\beta$ -D-glucopyranoside. (Table I).

Although the samples of both species were too small to isolate and identify the minor coumarins, the present results reveal nevertheless that the ability to synthesize trihydroxycoumarins must have existed in the ancestral stock of the Polytrichales before the breaking up of Pangaea.


### Experimental

The plant material was gathered in New Zealand by permission of the local authorities.

*Dendrologotrichum dendroides* (Hedw.) Broth. was collected in spring 1994 on the slopes of Mount Taranaki at 900 m s. m. by M. Veit; a voucher is kept in the private herbarium of H. G. (R 289) *Dawsonia superba* Grev. was collected by S. Bloor in January 1990 in Westland; a voucher is deposited at CHR (No. 465120).

Extraction and isolation were performed as described previously (Jung *et al.*, 1994 and 1995). By





1: R=CH<sub>3</sub>

2: R=glc

#### Acknowledgements

We are greatly indebted to Dr. S. Bloor (Industrial Research Ltd., Lower Hutt, New Zealand) and P. D. Dr. M. Veit (Universität Würzburg) for providing the plant material and to Dr. J. Zapp (Universität des Saarlandes) for running the NMR spectra.

Table I. <sup>13</sup>C- and <sup>1</sup>H-NMR data of **2** recorded in DMSO-d<sub>6</sub> at ambient temperature with 100 and 400 MHz respectively.

<sup>13</sup> C <sup>1</sup> H Position	<sup>13</sup> C	<sup>1</sup> H
2	159.9	-
3	111.4	6.24 d (9.6)
4	139.4	8.20 d (9.6)
5	145.8	-
6	98.7	6.85 s
7	148.2	-
8	128.3	-
9	142.5	-
10	104.7	-
1', 1''	100.9, 100.9	4.86 d (7.5), 4.82 d (7.4)
2', 2''	73.1, 73.1	3.74–3.10 m, 3.74–3.10 m
3', 3''	76.1, 76.8	3.74–3.10 m, 3.74–3.10 m
4', 4''	69.9, 69.9	3.74–3.10 m, 3.74–3.10 m
5', 5''	76.9, 76.9	3.74–3.10 m, 3.74–3.10 m
6', 6''	60.9, 60.9	3.74–3.10 m, 3.74–3.10 m

these methods 3 g branchlets of *D. dendroides* yielded 2 mg **1** and 38 g gametophytes of *D. su-perba* gave 7 mg **2**.

Jung M. (1993), Neue Tri- und Tetrahydroxycumarinderivate aus den Laubmoosen *Atrichum undulatum* (Hedw.) P. Beauv., *Polytrichum formosum* und *Tetraphis pellucida* (Hedw.), Dissertation, Saarbrücken.

Jung M., Zinsmeister H. D. and Geiger H. (1994), New three- and tetraoxygenated coumaringlucosides from the mosses *Atrichum undulatum* and *Polytrichum formosum*. Z. Naturforsch. **49c**, 697–702.

Jung M., Geiger H. and Zinsmeister H. D. (1995), Tri- and tetrahydroxycumarinderivatives from *Tetraphis pellucida*. Phytochemistry **39**, 379–381.

Raubuch J. (1998), Neue Cumarinderivate aus einigen Vertretern der Laubmoosklasse Bryopsida, Dissertation Saarbrücken.