

# Sesquiterpenoids from the Roots of *Solanum aethiopicum*

Hiroshi Nagase<sup>a</sup>, Toshinori Nagaoka<sup>b,\*</sup>, Ayako Watanabe<sup>a,d</sup>, Yoshiteru Sakata<sup>c</sup>  
and Teruhiko Yoshihara<sup>a,d</sup>

<sup>a</sup> Division of Applied Bioscience, Graduate School of Agriculture, Hokkaido University,  
North-9 West-9, Kita-ku, Sapporo 060–8589, Japan

<sup>b</sup> Faculty of Applied Biological Science, Hiroshima University, Kagamiyama 1-4-4, Higashi-  
Hiroshima 739–8528, Japan. Fax: +81-824-24-0791. E-mail: tnagaok@hiroshima-u.ac.jp

<sup>c</sup> National Research Institute of Vegetables, Ornamental Plants and Tea, Mii-machi 1823,  
Kurume 839–8503, Japan

<sup>d</sup> CREST, Japan Science and Technology Corporation (JST), Kawaguchi 332–0012, Japan

\* Author for correspondence and reprint requests

Z. Naturforsch. **56c**, 181–187 (2001); received December 2, 2000/January 22, 2001

*Solanum aethiopicum*, Roots, Sesquiterpenoid

Three new sesquiterpenoids, lubiminoic acid, epilubiminoic acid and aethione, and six known sesquiterpenoids, solavetivone, 3 $\beta$ -hydroxysolavetivone, 13-hydroxysolavetivone, anhydro- $\beta$ -rotunol, epilubimin and lubimin, were isolated from roots of *S. aethiopicum* L. Their structures were elucidated by spectroscopic data.