

# Lignan and Coumarin Glycosides from *Haplophyllum suaveolens*

Antoaneta Ivanova<sup>a</sup>, Bozhanka Mikhova<sup>a</sup>, Tatyana Stambolijska<sup>b</sup>  
and Ivanka Kostova<sup>a,\*</sup>

<sup>a</sup> Institute of Organic Chemistry with Centre of Phytochemistry and

<sup>b</sup> Institute of Physiology of Plants, Bulgarian Academy of Sciences, Sofia 1113, Bulgaria.  
E-mail: kostiv@bas.bg

\* Author for correspondence and reprint request

Z. Naturforsch. **56c**, 329–333 (2001); received December 28, 2000/February 6, 2001

*Haplophyllum suaveolens*, Dibenzylbutyrolactol, Arylnaphthalene Lignans,  
Coumarin Glycosides

Two new lignan glycosides of dibenzylbutyrolactol type (haplomarín) and aryl-naphthalene type (haploborín) were isolated from the aerial parts of *Haplophyllum suaveolens* together with the known aryl-naphthalene lignan glycoside arabelline and hydroxycoumarin glycoside xeroboside. So far arabelline is found only in *H. buxbaumii*. This is the first report of the isolation of xeroboside from *Haplophyllum*.