

# Iridoid and Phenylethanoid Glycosides from *Phlomis tuberosa* L.

Tayfun Ersöz<sup>a,\*</sup>, Stefanka Ivancheva<sup>b</sup>, Pınar Akbay<sup>c</sup>, Otto Sticher<sup>c</sup> and İhsan Çalış<sup>a</sup>

<sup>a</sup> Department of Pharmacognosy, Faculty of Pharmacy, Hacettepe University TR-06100, Ankara, Turkey.

Fax: +90-312-3114777. E-mail: tersoz@hacettepe.edu.tr

<sup>b</sup> Institute of Botany, Bulgarian Academy of Sciences, Sofia 1113, Bulgaria

<sup>c</sup> Department of Applied BioSciences, Institute of Pharmaceutical Sciences, ETH-Zurich, Winterthurerstr. 190, CH-8057 Zürich, Switzerland

\* Author for correspondence and reprint requests

Z. Naturforsch. **56c**, 695-698 (2001); received April 17/May 22, 2001

*Phlomis*, Iridoid Glucosides, Phenylethanoid Glycosides

A new iridoid glucoside, 8-*O*-acetylshanzhiside (**1**), was isolated from the aerial parts of *Phlomis tuberosa*, together with two known iridoid glucosides, shanzhiside methyl ester and lamalbide. The known phenylethanoid glycosides acteoside and forsythoside B were also obtained and characterized. The structure of **1** was determined by means of 1D- and 2D-NMR spectroscopic evidence.