

Andirol A and B, Two Unique 6-Hydroxymethylpterocarpenes from *Andira inermis*[§]

Carola Kraft^a, Kristina Jenett-Siems^{a,*}, Inga Köhler^a, Karsten Siems^b, Daniel Abbiw^c, Ulrich Bienzle^d and Eckart Eich^a

^a Institut für Pharmazie (Pharmazeutische Biologie), Freie Universität Berlin, Königin-Luise-Str. 2–4, D-14195 Berlin, Germany. Fax: +49 30 83 85 3729.
E-mail: kjsiems@zedat.fu-berlin.de

^b AnalytiCon Discovery GmbH, D-14473 Potsdam, Germany

^c Department of Botany, University Legon-Accra, Accra, Ghana

^d Institut für Tropenmedizin, Medizinische Fakultät Charité der Humboldt-Universität zu Berlin, D-14050 Berlin, Germany

* Author for correspondence and reprint requests

Z. Naturforsch. **57c**, 785–790 (2002); received May 10/May 31, 2002

Antiplasmodial Activity, Andirol A–B, Andinermol

From a methanolic extract of the leaves of *Andira inermis* (Fabaceae), andirol A and B, two compounds with a novel type of a rotenoid-related skeleton and andinermol, a new 2-aryl-3-hydroxymethyl-benzofuran could be isolated. Characterisation and structure elucidation of these compounds was achieved on the basis of their spectral data. In addition, the *in vitro* activities of the isolated compounds against both the chloroquine-sensitive strain PoW and the chloroquine-resistant clone Dd2 of *Plasmodium falciparum* have been evaluated.

Introduction

Andira inermis (W. Wright) H. B. K., Fabaceae, a plant remedy native from Mexico to Southern America also occurs often as an ornamental in West Tropical Africa. Recently, we reported the isolation of 2-arylbenzofuran-3-carbaldehydes (andinermals) from the leaves of a Panamanian sample. These, as well as the also isolated isoflavones calycosin and genistein proved to be active against both the chloroquine-sensitive strain PoW and the chloroquine-resistant clone Dd2 of *Plasmodium falciparum* with IC₅₀ values between 2.0 µg/ml/7.4 µM to 9.8 µg/ml/34.5 µM (Kraft *et al.*, 2001). In this study we investigated the leaves of this species collected in Ghana and obtained two novel rotenoid-type flavonoids and one new 2-aryl-3-hydroxymethyl-benzofuran. Furthermore, the antiplasmodial effects of the isolated compounds have been assayed.


Results and Discussion

In addition to the already known constituents of *A. inermis* (Kraft *et al.*, 2000, 2001), three novel

compounds could be obtained by bioassay-guided fractionation of the CHCl₃ fraction. Their structural elucidation was carried out by EIMS, ¹H NMR, ¹³C NMR, HMBC, as well as NOE experiments.


Compound **1** (Fig. 1) gave a molecular ion peak at *m/z* 342 in the EIMS spectrum, corresponding to a molecular formula of C₁₈H₁₄O₇. The ¹H NMR spectrum (Table I) indicated the presence of a methylendioxy group (6.01 ppm, s, 2 H), one isolated aromatic proton (6.81 ppm, s, 1 H) and one methoxy group (4.16 ppm, s, 3 H). Additionally the ¹H NMR displayed characteristic signals of a 1,2,4-trisubstituted aromatic ring and the typical pattern of an ABX-system (4.03 ppm, dd, 1H; 4.46 ppm, dd, 1H; 5.18 ppm, dd, 1H) similar to those described for 12a-hydroxyrotenoids (Abe *et al.*, 1985). In the HMBC (Table I) the protons at C-6 and C-6a showed correlations to a carbon at 117.9 ppm corresponding to a double bond. In addition, H-6a correlated to a carbon at 150.2 ppm. The ¹³C NMR chemical shifts of the double bond between C-12 and C-12a at δ 150.2 ppm and 117.9 ppm, respectively, were characteristic for an enolether. On the other hand, the ring between C-6 and C-4a which is typical for rotenoids has been opened. Thus, compound **1** represented a re-

[§] Part 7 in the series “Herbal remedies traditionally used against malaria”. For part 6 see Onogi *et al.*




1 R = OCH₃
2 R = H

numbering system based on the rotenoids




1 R = OCH₃
2 R = H

numbering system based on the pterocarpanes



3



4

Fig. 1. Compounds isolated from the leaves of *Andira inermis*.

arranged rotenoid derivative (the numbering system of the rotenoids has been kept, Fig. 1). The positions of the methylendioxy group and of the hydroxy group could be deduced from the HMBC, whereas the position of the methoxy group at C-4a could be determined by NOE experiments: irradiation of the singlet at 4.16 ppm led to the enhancement of the H-6a signal (5.18 ppm). Compound **2** showed similar signals in the ¹H NMR, its molecular formula was determined as C₁₇H₁₂O₆. In contrast to **1** no methoxy group could be observed. The singlets at 7.08 and 7.14 ppm represented the *para*-positioned protons at C-2 and C-4a, respectively. The shifts and coupling patterns of the other aromatic signals and the methylendioxy group were nearly identical to those of **1**. Thus, **2** is the 4a-demethoxy derivative of **1**; for these com-

pounds we propose the names andiro A (**1**) and B (**2**).

The molecular formula of **3** was calculated as C₁₉H₂₀O₇ by HRMS. The ¹H NMR displayed characteristic signals of a pair of *meta*-coupled protons at 6.38 ppm (d, J = 1.9 Hz, H-5) and 6.69 ppm (d, J = 1.9 Hz, H-7), as well as of two *ortho*-coupled protons at 6.75 ppm (d, J = 8.5 Hz, H-5') and 6.98 ppm (d, J = 8.5 Hz, H-6') and four methoxy groups. Furthermore, one hydroxymethyl moiety could be characterized in the ¹H-NMR by a doublet at 4.71 ppm (J = 7.0 Hz, 2H) and by a downfield signal at 56.5 ppm in the ¹³C NMR. The spectral data were comparable to those of 2-(3',4'-dimethoxyphenyl)-3-hydroxymethyl-7-methoxybenzofuran which could be prepared by reduction of the corresponding 3-formyl derivative by Stom-

Table I. ^1H NMR data [400 MHz, CD_3COCD_3 , δ_{H} (ppm), J (Hz)], ^{13}C -NMR data [100.6 MHz, CD_3COCD_3 , δ_{C} (ppm)] and HMBC correlations of andirolo A (**1**) and ^1H NMR data [400 MHz, CD_3COCD_3 , δ_{H} (ppm), J (Hz)] of andirolo B (**2**).

Proton ^a	Carbon ^a	Compound 1			Compound 2 δ_{H}
		δ_{H}	δ_{C}	HMBC Correlations ^e	
—	C-1	—	148.4 ^b		—
H-2	C-2	6.81, 1 H, s	136.7		7.08, 1 H, s
—	C-3	—	133.5		—
—	C-4	—	148.9 ^b		—
H-4a	C-4a	—	88.8	1, 3, 4, 12a, 12b	7.14, 1 H, s
H-6	C-6	4.03, 1 H, dd (1.9 and 12.4)	75.1	6, 12a	4.12, 1 H, dd (2.2 and 12.2)
H-6'		4.46, 1 H, dd (3.8 and 12.4)			4.41, 1 H, dd (4.4 and 12.2)
H-6a	C-6a	5.18, 1 H, dd (1.9 and 3.8)	65.9	12, 12a	5.06, 1 H, m
—	C-7a	—	159.5 ^c		—
H-8	C-8	6.60, 1 H, d (2.3)	108.2	7a, 9, 10, 11a	6.60, 1 H, d (2.3)
—	C-9	—	159.8 ^c		—
H-10	C-10	6.71, 1H, dd (2.3 and 8.6)	111.8	11a	6.72, 1H, dd (2.3 and 8.8)
H-11	C-11	7.76, 1 H, d (8.6)	127.8	7a, 9, 12	7.76, 1 H, d (8.8)
—	C-11a	—	114.2		—
—	C-12	—	150.2		—
—	C-12a	—	117.9 ^d		—
—	C-12b	—	116.5 ^d		—
4a-OCH ₃	OCH ₃	4.16, 3 H, s	60.6	2	—
-O-CH ₂ -O-	-O-CH ₂ -O-	6.01, 2 H, s	102.2	3, 4	6.03, 2 H, s

^a Numbering system based on the rotenoids; ^{b,c,d} Interchangeable; ^e CD_3OD as solvent.

berg *et al.* (1997). In addition they were similar to andinermal A (**4**), a 2-aryl-3-carbaldehyde, previously isolated from *A. inermis* (Kraft *et al.*, 2001). Thus, **3** represented the reduced form of **4** with an additional methyl group. The position of the three methoxy groups at 3.86, 3.95, and 3.98 ppm (6-OCH₃, 4'-OCH₃, 4-OCH₃, respectively) were established by NOE experiments whereas the ^{13}C NMR chemical shift of the fourth methoxy group at 61.13 ppm pointed to the C-2' position in ring B as in **4**. This could be confirmed by NOE irradiation of the doublet at 4.71 ppm which led to the enhancement of the H-6' signal (6.98 ppm). **3** represents a novel compound, which we named andinermol.

In the *in vitro* antiplasmoidal assay against *Plasmodium falciparum* the methanolic extract exhibited no activity. After successive extraction we found an enhanced activity in the CHCl₃ fraction (IC₅₀: 10.6 $\mu\text{g}/\text{ml}$ [Dd2 clone of *P. falciparum*]). Although isolated from the most active fractions, andirolo A (**1**) showed no remarkable antiplas-

modial activity (IC₅₀ values: 15.0 $\mu\text{g}/\text{ml}$ /43.9 μM [PoW strain of *P. falciparum*]; 42.5 $\mu\text{g}/\text{ml}$ /124.3 μM [Dd2]). Andinermol (**3**) exhibited slightly lower IC₅₀ values (8.6 $\mu\text{g}/\text{ml}$ /23.9 μM [PoW]; 13.7 $\mu\text{g}/\text{ml}$ /38.1 μM [Dd2]), but it was less active than the related carbaldehyde andinermal A (**4**; IC₅₀ values: 2.1 $\mu\text{g}/\text{ml}$ /6.1 μM [PoW]; 3.9 $\mu\text{g}/\text{ml}$ /11.3 μM [Dd2]; Kraft *et al.*, 2001). Due to the lack of sufficient amounts of andirolo B (**2**), it could not be tested.

In conclusion, during our investigation on the phytochemical spectrum of *A. inermis*, we could isolate from the leaves two compounds (**1**, **2**) with a unique heterocyclic structure. In contrast to the 2-arylbenzofuran-3-carbaldehydes they exhibited no antiplasmoidal activity. Their carbon skeleton seems to be related to that of the rotenoids. Rotenoids are a class of isoflavonoids characterized by the presence of an extra carbon atom in an additional heterocyclic ring. This system originates in nature by oxidative cyclization of a 2'-methoxy-isoflavone (Harborne, 1988). In the case of **1** and

2, the cyclization must be due to the enolic hydroxyl group in position 12 and a hydroxy group in position 1. Up to now, no similar compounds are described in literature.

Whereas several isoflavones (e.g. genistein, formononetin, and calycosin) and the 2-arylbenzofuran-3-carbaldehyde andinermal A (**4**) as well as andiro A (**1**) could be detected in both the Panamanian plant material and the leaves harvested in Ghana, andiro B (**2**) and andinermol (**3**), a 2-aryl-3-hydroxymethyl-benzofuran could only be found in the African plant. 2-Aryl-benzofurans are normally derived from flavonoids or pterocarpans. They seem to occur rarely in nature. From literature simple benzofurans like 2-(2,4-dihydroxyphe-nyl)-5,6-methylendioxy-benzofuran from *Artemisia indica* Willd., Asteraceae, (Chanphen *et al.*, 1998) or ebenfuran I from *Onobrychis ebenoides* Bioss & Spuner, Fabaceae (Halabalaki *et al.*, 2000) are known, as well as 3-methyl-benzofurans like parvifuran from *Dalbergia parviflora* Prain, Fabaceae, (Muangnoicharoen and Frahm, 1981), benzofuran-3-carbaldehydes like ebenfuran II and III (Halabalaki *et al.*, 2000) or andinermal A-C (Kraft *et al.*, 2001), and benzofuran-3-carboxylic acids like norwedelic acid from *Wedelia calendulacea* Less., Asteraceae (Govindachari and Premila, 1985). In contrast, **3** represents the first natural 2-aryl-3-hydroxymethyl-benzofuran. Up to now, such structural feature was only described in semi-synthetic reduction products from the related carbaldehydes (Stomberg *et al.*, 1997) or as synthetic educts in chemical syntheses (Fukui *et al.*, 1969).

Experimental

General

For fractionation silica gel 60 (63–200 µm) was used. Preparative HPLC was performed on a Knauer instrument equipped with Eurochrom 2000 on a Nucleosil P 300 C-18 (10 µm) column and medium pressure liquid chromatography (MPLC) on a LiChroprep® RP-18 (40–63 µm) column. EIMS were recorded on a Finnigan MAT CH7A (70 eV); FAB MS and HR-EIMS on a Finnigan MAT 711 (80 eV). ¹H NMR, ¹³C NMR, NOE, and HMBC spectra were obtained on a Bruker AVANCE DPX 400 MHz or a Bruker

DRX 500 MHz spectrometer (TMS as int. standard; CD₃COCD₃ or CDCl₃ as solvent).

Plant material

The leaves of *A. inermis* were collected in the Upper East Region of Ghana during March 2000. Voucher specimens (GC 47682) were identified by Dr. D. Abbiw and deposited at the Department of Botany, University Legon-Accra, Ghana.

Extraction and isolation

Ground leaves (400 g) were extracted four times with 1.5 l MeOH-H₂O (8:2, v/v) for 24 h at room temperature. MeOH was evaporated and the remaining aqueous phase (ca. 500 ml) was successively extracted with petrol ether, CHCl₃, and EtOAc (four times with 300 ml each) to yield four fractions, which were tested against *P. falciparum*. The CHCl₃ fraction proved to be most active.

The oily residue (3.4 g) was subjected to column chromatography on silica gel 60 (50 g) and eluted with CHCl₃, CHCl₃-MeOH mixtures, and MeOH to yield seven fractions. Due to the results of the bioassay, fraction 3 (190 mg, eluted with CHCl₃-MeOH 9:1 [v/v]; IC₅₀: 4.2 µg/ml [PoW]) and fraction 4 (420 mg, eluted with CHCl₃-MeOH 8:2 [v/v]; IC₅₀: 8.0 µg/ml [PoW]) were further separated. From fraction 3 we gained three main peaks by preparative HPLC (RP-18) with a gradient of MeOH-H₂O 3:7 (v/v) to MeOH-H₂O 7:3 (v/v) in 40 min at a flow rate of 5.5 ml/min, which were purified by preparative TLC. Peak 1 and 2 were purified on TLC (cyclohexane-EtOAc 3:7 [v/v]) and yielded **3** (R_f: 0.35; 1 mg) and **1** (R_f: 0.37; 6.3 mg), respectively. From peak 3 compound **4** could be identified by comparison of spectral data (Kraft *et al.*, 2001). The purification of fraction 4 was performed by MPLC with MeOH-H₂O mixtures. Again the fourth fraction (350 mg, eluted with MeOH-H₂O 7:3 [v/v]; IC₅₀: 4.7 µg/ml [PoW]) was the most active one. After a second column chromatography on silica gel (10 g) with CHCl₃, CHCl₃-MeOH mixtures and MeOH we gained again **1** (2 mg; eluted with CHCl₃-MeOH 99:1 [v/v]). By preparative TLC (CHCl₃-MeOH 95:5 [v/v]; developed twice) of the fraction eluted with CHCl₃-MeOH 99:1 and 95:5 [v/v], **2** could be isolated (R_f: 0.46; < 1 mg).

Spectroscopic data

3-Hydroxy-7-methoxy-8,9-methylendioxy-6-hydroxymethylpterocarpene (**1**, andirol A): Yellow needles; $[\alpha]_D^{20}$: + 5.33° (CHCl₃, c = 0.075); EIMS *m/z* (rel. Int.): 342 [M]⁺ (92), 324 [M-H₂O]⁺ (100), 313, (53), 309, (40); (+)-FABMS: *m/z*: 342 [M]⁺, 325 [M-H₂O+H]⁺; (-)-FABMS: *m/z*: 341 [M-H]⁻; HRMS *m/z*: 342.07368 (C₁₈H₁₄O₇, calculated for 342.07396), 324.06315 (C₁₈H₁₂O₆, calculated for 324.06339); ¹H NMR and ¹³C-NMR: see Table I.

3-Hydroxy-8,9-methylendioxy-6-hydroxymethylpterocarpene (**2**, andirol B): Yellow needles; EIMS *m/z* (rel. Int.): 312 [M]⁺ (31), 294 [M-H₂O]⁺ (43), 167 (20), 149 (49), 28 (100); HRMS *m/z*: 312.06352 (C₁₇H₁₂O₆, calculated for 312.063390), 294.05271 (C₁₇H₁₀O₅, calculated for 294.05283); ¹H NMR: see Table I.

2-[2',4'-Dimethoxy-3'-hydroxyphenyl]-4,6-dimethoxy-3-hydroxymethyl-benzofuran (**3**, andinermol): Yellow needles; EIMS *m/z* (rel. Int.): 360 [M]⁺ (100), 342 [M-H₂O]⁺ (10); HRMS *m/z*: 360.12078 (C₁₉H₂₀O₇, calculated for 360.12091); ¹H NMR (400 MHz, CDCl₃) δ 6.98 (1H, d, *J* = 8.5 Hz, H-6'), 6.75 (1H, d, *J* = 8.5 Hz, H-5'), 6.69 (1H, d, *J* = 1.9 Hz, H-7), 6.38 (1H, d, *J* = 1.9 Hz, H-5), 5.95 (1H, brs, 3'-OH), 4.71 (2H, d, *J* = 7.0 Hz, H_{a,b}-8), 3.98 (3H, s, 4-OCH₃), 3.95 (3H, s, 4'-OCH₃), 3.86 (3H, s, 6-OCH₃), 3.71 (3H, s, 2'-OCH₃), 3.24 (1H, t, *J* = 7.0 Hz, 8-OH); ¹³C NMR (100.6 MHz, CDCl₃) δ 159.8, 159.2, 156.5 (C-4, C-6, C-7a), 153.2 (C-4'), 148.5 (C-2'), 138.7 (C-3'),

121.7 (C-6'), 116.6 (C-3), 108.3, 106.8 (C-3a, C-5'), 94.3 (C-5), 88.7 (C-7), 61.1 (2'-OCH₃), 56.5 (C-8), 56.3, 55.8, 55.8 (4-OCH₃, 6-OCH₃, 4'-OCH₃), n. d. (C-2, C-1').

In vitro antiplasmodial activity

The bioassay was performed as previously described (Kraft *et al.*, 2000). *P. falciparum* strain PoW (IC₅₀ for chloroquine = 0.008 µg/ml/0.015 µM) and the clone Dd2 (IC₅₀ = 0.092 µg/ml/0.18 µM) were maintained in continuous culture in human red blood cells (A⁺) as previously described (Trager and Jensen, 1976). Substances were dissolved in DMSO (20 mg/ml) and diluted in medium to final concentrations between 50 and 0.78 µg/ml. The antiplasmodial assay, according to Desjardins *et al.* (1979) was analysed by means of the microculture radioisotope technique after 42 h.

Acknowledgements

The authors are indebted to Mrs U. Ostwald, Institut für Organische Chemie, FU Berlin, for providing the HR-EIMS and FAB spectra and to the colleagues at the Department of Botany, University Legon-Accra for their support in collecting the plant material. This study was supported by a grant of the Deutsche Pharmazeutische Gesellschaft to K. J.-S. and from the Humboldt-Universität zu Berlin (N-1/01/01) to I. K.

- Abe F., Donelly D. M. X., Moretti C. and Polonsky J. (1985), Isoflavonoid constituents from *Dalbergia monnieria*. *Phytochemistry* **24**, 1071–1076.
- Chanphen R., Thebtaranonth Y., Wanauppathamkul S. and Yuthavong Y. (1998), Antimalarial principles from *Artemisia indica*. *J. Nat. Prod.* **61**, 1146–1147.
- Desjardins R. E., Canfield C. J., Haynes J. D. and Chulay J. D. (1979), Quantitative assessment of antimalarial activity *in vitro* by a semiautomated microdilution technique. *Antimicrob. Agents Chemother.* **16**, 710–718.
- Fukui K., Nakayama M. and Tsuzuki K. (1969), Synthesis of pterocarpan. IV. A new synthesis of 3,8,9-trimethoxypterocarpan. *Bull. Chem. Soc. Jap.* **42**, 2395–2396.
- Govindachari T. R. and Premila M. S. (1985), The benzofuran norweddelic acid from *Wedelia calendulacea*. *Phytochemistry* **24**, 3068–3069.
- Halabalaki M., Aligiannis N., Papoutsi Z., Mitakou S., Moutsatsou P., Sekeris C. and Skaltsounis A. L. (2000), Three new arylbenzofurans from *Onobrychis ebenoides* and evaluation of their binding affinity for the estrogen receptor. *J. Nat. Prod.* **63**, 1672–1674.
- Harborne J. B. (1988), The Flavonoids: Advances in Research Since 1980. Chapman and Hall, London, pp. 152–157.
- Kraft C., Jenett-Siems K., Siems K., Gupta M. P., Bienzle U. and Eich E. (2000) Antiplasmodial activity of isoflavones from *Andira inermis*. *J. Ethnopharmacol.* **73**, 131–135.
- Kraft C., Jenett-Siems K., Siems K., Solis P. N., Gupta M. P., Bienzle U. and Eich E. (2001), Andinermals A–C, antiplasmodial constituents from *Andira inermis*. *Phytochemistry* **58**, 769–774.
- Muangnoicharoen N. and Frahm A. W. (1981), Arylbenzofurans from *Dalbergia parviflora*. *Phytochemistry* **20**, 291–293.
- Onogi B., Kraft C., Köhler I., Freund M., Jenett-Siems K., Siems K., Beyer G., Melzig M. F., Bienzle U. and Eich E. (2002), Antiplasmodial activity of naphthoquinones and one anthraquinone from *Stereospermum kunthianum*. *Phytochemistry* **60**, 39–44.
- Stomberg R., Li S., Lundquist K. and Wallis A. F. A. (1997), Crystal structure of 2-(3,4-dimethoxyphenyl)-3-hydroxymethyl-7-methoxybenzo[b]furan, $C_{18} H_{18} O_5$. *Z. Kristallogr. – New Cryst. Struct.* **212**, 473–474.
- Trager W. and Jensen J. B. (1976), Human malaria parasites in continuous culture. *Science* **193**, 673–675.